

**ACIP-Recommended Haemophilus influenzae type b (Hib)  
Routine Vaccine Schedule**

Type	Vaccine	2 months	4 months	6 months	12-15 months
PRP-T	ActHIB	X (1st)	X (2nd)	X (3rd)	X
	Pentacel*	X (1st)	X (2nd)	X (3rd)	X
	Hiberix	X (1st)	X (2nd)	X (3rd)	X
PRP-OMP	PedvaxHIB	X (1st)	X (2nd)	—	X
	Vaxelis	X (1st)	X (2nd)	X (3rd)	X

\*The recommended age for the 4<sup>th</sup> dose of Pentacel is 15-18 months, but it can be given as early as 12 months, provided at least 6 months have elapsed since the 3<sup>rd</sup> dose.

**Haemophilus influenzae type b Vaccine Detailed Schedule for  
Unvaccinated Children**

Vaccine	Age at 1st Dose (months)	Primary series	Booster
PRP-T	2-6 months	3 doses, 8 weeks apart	12-15 months
	7-11 months	2 doses, 4 weeks apart	12-15 months
	12-14 months	1 dose	2 months later
	15-59 months	1 dose	--
PRP-OMP	2-6 months	2 doses, 8 weeks apart	12-15 months
	7-11 months	2 doses, 4 weeks apart	12-15 months
	12-14 months	1 dose	2 months later
	15-59 months	1 dose	--